

Closed System of Equations on a Brane

Yu. V. Shtanov*

Bogolyubov Institute for Theoretical Physics, Kiev 03143, Ukraine

(February 1, 2008)

We obtain a generic closed system of equations on a brane that describes its inner evolution and give a method for extending solutions on the brane to the bulk. We also discuss the cosmological implications of the closed system of equations obtained. We consider bulk spaces with both spacelike and timelike extra dimension, with and without the Z_2 symmetry of reflection relative to the brane.

The idea that our four-dimensional world can be described as a timelike hypersurface (brane) embedded in or bounding a five-dimensional manifold has been extensively studied already for several years. In its ideal formulation, it is only the evolution on the brane that is observable by experimenters themselves residing in the four-dimensional braneworld, and the role of the five-dimensional bulk is to provide a special form of the laws that govern this evolution on the brane, for example, the unusual form of the law of gravity on relatively small scales, as was observed in the seminal papers [1].

In solving equations for the brane, there naturally arises the question about the additional conditions that must be satisfied by the solution in the bulk. Usually, one considers another brane or system of branes on which one can impose certain boundary conditions and require that the solution be nonsingular in the space between the branes. Very often, however, no such additional conditions in the bulk are actually imposed while solving field equations in the neighborhood of the brane of interest. Such an approach has some right to exist since, as was pointed out, the only observable evolution is the evolution on a single braneworld.

*E-mail: shtanov@bitp.kiev.ua

In this letter, we would like to point out that, *in this approach*, and in the absence of matter fields in the bulk, all the information about the evolution on the brane is contained in a system of closed equations [Eqs. (20)–(23) below], without the necessity of solving field equations in the bulk. A similar observation in the cosmological context was previously made in [2] and recently noted in [3] for the linearized equations in the case of Z_2 symmetry. In [4], it was argued that cosmological density and curvature perturbations on sufficiently large spatial scales are described by a closed system of equations on the brane. Here, we would like to point out that this feature is rather general.

We also describe a method for constructing the extension of solution on the brane to the bulk using Gaussian normal coordinates in the neighbourhood of the brane.

In most of the braneworld models, the extra dimension is assumed to be spacelike, so that the brane is embedded in a Lorentzian five-dimensional manifold. However, no physical principle appears to prevent us from considering a complementary model in which the extra dimension is timelike, so that the brane is a Lorentzian boundary of a five-dimensional space with signature $(-, -, +, +, +)$. Such general embeddings in the case of Z_2 symmetry were considered in [5] and also in [6]. Models with both signatures of the five-dimensional bulk space will be examined in this paper.

To begin with, we consider a particular case where a four-dimensional hypersurface (brane) $\mathcal{B}$ is the boundary of a five-dimensional Riemannian manifold (bulk) $\mathcal{M}$ with non-degenerate Lorentzian induced metric. The action of the theory has the natural general form

$$S = M^3 \left[\int_{\mathcal{M}} (\mathcal{R} - 2\Lambda) - 2\epsilon \int_{\mathcal{B}} K \right] + \int_{\mathcal{M}} L_5(g_{ab}, \Phi) + \int_{\mathcal{B}} (m^2 R - 2\sigma) + \int_{\mathcal{B}} L_4(h_{ab}, \phi). \quad (1)$$

Here, $\mathcal{R}$ is the scalar curvature of the five-dimensional metric g_{ab} on $\mathcal{M}$, and R is the scalar curvature of the induced metric $h_{ab} = g_{ab} - n_a n_b$ on $\mathcal{B}$, where n^a is the vector field of the *inner* unit normal to $\mathcal{B}$. The quantity $K = K_{ab} h^{ab}$ is the trace of the symmetric tensor of extrinsic curvature $K_{ab} = h^c_a \nabla_c n_b$ of $\mathcal{B}$ in $\mathcal{M}$. The parameter $\epsilon = 1$ if the signature of the bulk space is Lorentzian, so that the extra dimension is spacelike, and $\epsilon = -1$ if the

signature is $(-, -, +, +, +)$, so that the extra dimension is timelike. The symbols $L_5(g_{ab}, \Phi)$ and $L_4(h_{ab}, \phi)$ denote, respectively, the Lagrangian densities of the five-dimensional matter fields Φ and four-dimensional matter fields ϕ the dynamics of which is restricted to the brane $\mathcal{B}$ so that they interact only with the induced metric h_{ab} . Some of the fields ϕ , in principle, may represent restrictions of some of the fields Φ to the boundary $\mathcal{B}$. All integrations over $\mathcal{M}$ and over $\mathcal{B}$ are taken with the natural volume elements $\sqrt{-\epsilon g} d^5x$ and $\sqrt{-h} d^4x$, respectively, where g and h are the determinants of the matrices of components of the metric on $\mathcal{M}$ and of the induced metric on $\mathcal{B}$, respectively, in coordinate basis. The symbols M and m denote the Planck masses of the corresponding spaces, Λ is the five-dimensional cosmological constant, and σ is the brane tension.

In this paper, we systematically use the notation and conventions of [7]. In particular, we use the one-to-one correspondence between tensors in $\mathcal{B}$ and tensors in $\mathcal{M}$ which are invariant under projection to the tangent space to $\mathcal{B}$, i.e., tensors $T^{a_1 \dots a_k}_{b_1 \dots b_l}$ such that

$$T^{a_1 \dots a_k}_{b_1 \dots b_l} = h^{a_1}_{c_1} \dots h^{a_k}_{c_k} h_{b_1}^{d_1} \dots h_{b_l}^{d_l} T^{c_1 \dots c_k}_{d_1 \dots d_l}. \quad (2)$$

The term containing $m^2 R$ is sometimes missing from the action, or its contribution is missing from the equations of motion. However, in general, this term is essential since it is generated as a quantum correction to the matter action in (1); in this paper, we take it into account following [8–10].

Variation of action (1) gives the equation of motion in the five-dimensional bulk:

$$\mathcal{G}_{ab} + \Lambda g_{ab} = \frac{1}{M^3} T_{ab}, \quad (3)$$

and on the brane $\mathcal{B}$:

$$m^2 G_{ab} + \sigma h_{ab} = \epsilon M^3 S_{ab} + \tau_{ab}, \quad (4)$$

where $\mathcal{G}_{ab}$ and G_{ab} are the Einstein's tensors of the corresponding spaces, $S_{ab} \equiv K_{ab} - K h_{ab}$, and T_{ab} and τ_{ab} denote the five-dimensional and four-dimensional stress–energy tensors of

matter, respectively. It is the presence of the tensor S_{ab} in the equation of motion (4) that makes the inner dynamics on the brane $\mathcal{B}$ unusual.¹

The Codazzi relation

$$D_a S^a_b = \mathcal{R}_{cd} n^d h^c_b, \quad (5)$$

where D_a is the (unique) derivative on the brane $\mathcal{B}$ associated with the induced metric h_{ab} , together with Eq. (4) and bulk equation (3) imply the relation

$$D_a \tau^a_b = -\epsilon T_{cd} n^d h^c_b. \quad (6)$$

Thus, the four-dimensional stress–energy tensor is covariantly conserved on the brane if and only if the right-hand side of (6) is vanishing at the brane, in particular, if the five-dimensional stress–energy tensor T_{ab} is a linear combination of g_{ab} and h_{ab} at the brane. In this paper, we assume this to be the case by setting $T_{ab} = 0$, i.e., by considering only vacuum Einstein equation in the bulk.

Equation (4) involves the tensor S_{ab} which is constructed from the tensor of extrinsic curvature of the brane, so this equation is not closed with respect to the intrinsic evolution on the brane. However, it is possible to obtain an equation which involves only four-dimensional fields on the brane. In doing this, we follow the procedure of [11,12]. We use the Gauss relation on the brane:

$$R_{abc}{}^d = h_a{}^f h_b{}^g h_c{}^k h^d{}_j \mathcal{R}_{fgk}{}^j + \epsilon \left(K_{ac} K_b{}^d - K_{bc} K_a{}^d \right). \quad (7)$$

Contracting this relation and taking into account Eq. (3) with $T_{ab} = 0$, one obtains the equation

$$\epsilon (R - 2\Lambda) + K_{ab} K^{ab} - K^2 \equiv \epsilon (R - 2\Lambda) + S_{ab} S^{ab} - \frac{1}{3} S^2 = 0, \quad (8)$$

which we also expressed in terms of $S_{ab} = K_{ab} - h_{ab} K$ and $S = h^{ab} S_{ab}$. This is the well-known constraint equation on the brane from the viewpoint of the gravitational dynamics in

¹Those interested in the derivation of (4) may look into the appendix of [10].

the five-dimensional bulk. Now we express the tensor S_{ab} in terms of the four-dimensional quantities using Eq. (4) and substitute it into Eq. (8). We obtain

$$S_{ab} = \frac{\epsilon}{M^3} \left(m^2 G_{ab} + \sigma h_{ab} - \tau_{ab} \right), \quad (9)$$

$$\epsilon M^6 (R - 2\Lambda) + \left(m^2 G_{ab} + \sigma h_{ab} - \tau_{ab} \right) \left(m^2 G^{ab} + \sigma h^{ab} - \tau^{ab} \right) - \frac{1}{3} \left(m^2 R - 4\sigma + \tau \right)^2 = 0, \quad (10)$$

where $\tau = h^{ab} \tau_{ab}$.

Together with the equations for other fields on the brane, Eq. (10) forms a system of closed equations on the brane to be solved. Note that the five-dimensional part of the curvature tensor in the bulk (the projected Weyl tensor) drops from this equation because the complete trace of the Gauss relation (7) has been taken. Once its solution is obtained, one can calculate the tensor of extrinsic curvature using Eq. (9) and then, if necessary, solve the bulk equations in the neighborhood of the brane with the given boundary data on the brane.

One way of doing this consists in choosing the Gaussian normal coordinates in the bulk generated by the Minkowski coordinates x^α , $\alpha = 0, \dots, 3$, on the brane, so that the metric of the solution in these coordinates has the form

$$ds_5^2 = \epsilon dy^2 + h_{\alpha\beta}(y, x) dx^\alpha dx^\beta, \quad (11)$$

where $y \geq 0$ is the fifth coordinate in the bulk, and the brane corresponds to $y = 0$. Introducing also the tensor of extrinsic curvature with components $K_{\alpha\beta}$ on every hypersurface $y = \text{const}$ in the bulk, one can obtain the following system of differential equations for the components $h_{\alpha\beta}$ and K^α_β :

$$\begin{aligned} \frac{\partial K^\alpha_\beta}{\partial y} &= \epsilon \left[R^\alpha_\beta - \frac{1}{6} (R + 2\Lambda) \delta^\alpha_\beta \right] - K K^\alpha_\beta - \frac{1}{6} \left(K^\mu_\nu K^\nu_\mu - K^2 \right) \delta^\alpha_\beta \\ &= \epsilon \left(R^\alpha_\beta - \frac{2}{3} \Lambda \delta^\alpha_\beta \right) - K K^\alpha_\beta, \end{aligned} \quad (12)$$

$$\frac{\partial h_{\alpha\beta}}{\partial y} = 2h_{\alpha\gamma} K^\gamma_\beta, \quad (13)$$

where R^α_β are the components of the Ricci tensor of the metric $h_{\alpha\beta}$ induced on the hypersurface $y = \text{const}$, $R = R^\alpha_\alpha$ is its scalar curvature, and $K = K^\alpha_\alpha$ is the trace of the tensor of extrinsic curvature. The second equality in (12) is true by virtue of the constraint equation (8). Equations (12) and (13) together with the constraint equation (8) represent the 4+1 splitting of the Einstein equations in the Gaussian normal coordinates. Solution of Eqs. (12) and (13) in the bulk always exists in some neighborhood of the brane.

We illustrate this method on the example of the generalization of the Randall–Sundrum (RS) solution [1] to the case of arbitrary signature of the fifth dimension and arbitrary Ricci-flat brane. In this solution, the matter on the brane is in the vacuum state, and the induced metric on the brane is Ricci-flat, so that $\tau_{\alpha\beta} = 0$ and $R^\alpha_\beta = 0$ at the brane. It follows from Eq. (4) that $K^\alpha_\beta \propto \delta^\alpha_\beta$ at the brane. Then Eqs. (12) and (13) imply that $K^\alpha_\beta(y, x) \equiv \frac{1}{4}K(y)\delta^\alpha_\beta$ and the induced metric $h_{\alpha\beta}(y, x)$ is Ricci-flat on every hypersurface $y = \text{const}$, so that $R^\alpha_\beta(y, x) \equiv 0$. After this, Eqs. (8) and (4) give

$$K^2 = -\frac{8}{3}\epsilon\Lambda, \quad K = -\frac{4\epsilon\sigma}{3M^3} \quad \Longrightarrow \quad \epsilon\Lambda = -\frac{2\sigma^2}{3M^6}, \quad (14)$$

which, in particular, means that $\epsilon\Lambda$ must be negative. Now Eq. (13) can easily be solved, and the resulting metric is

$$ds^2 = \epsilon dy^2 + \exp\left(-\frac{\epsilon\sigma}{3M^3}y\right) h_{\alpha\beta}(x) dx^\alpha dx^\beta, \quad (15)$$

where $h_{\alpha\beta}(x)$ are the components of the Ricci-flat metric on the brane.

If one is interested in the evolution on the brane only, and if no additional boundary and regularity conditions are specified for the bulk, then it suffices to solve only Eq. (10) together with the matter equations on the brane. As an example, let us show that the cosmological equations for the brane follow immediately from Eq. (10). We consider a homogeneous and isotropic cosmological model with the cosmological time t , scale factor $a(t)$, energy density $\rho(t)$, and stress $p(t)$. First, using the stress–energy conservation equation, we obtain

$$M^6 a^3 \dot{a} \left(S_{ab} S^{ab} - \frac{1}{3} S^2 \right) = -\frac{1}{3} \frac{d}{dt} \left[3m^2 (\dot{a}^2 + \kappa) - a^2 (\rho + \sigma) \right]^2, \quad (16)$$

where $\kappa = 0, \pm 1$ indicates the spatial curvature of the model. The expression $a^3 \dot{a}(R - 2\Lambda)$ is also a total derivative; hence, Eq. (10) can be integrated once with the result (obtained by different methods in [8,13] for the case of $\epsilon = 1$)

$$m^4 \left(H^2 + \frac{\kappa}{a^2} - \frac{\rho + \sigma}{3m^2} \right)^2 = \epsilon M^6 \left(H^2 + \frac{\kappa}{a^2} - \frac{\Lambda}{6} - \frac{C}{a^4} \right), \quad (17)$$

where C is the integration constant and $H \equiv \dot{a}/a$. For $m = 0$, this equation passes to a generalization of the well-known result [11,2]:

$$H^2 + \frac{\kappa}{a^2} = \frac{\Lambda}{6} + \frac{C}{a^4} + \epsilon \frac{(\rho + \sigma)^2}{9M^6}. \quad (18)$$

Some consequences of this equation for $\epsilon = -1$ are discussed in [6]. For $M = 0$, Eq. (17) becomes the standard cosmological equation in Einstein's theory.

Thus far, we considered the case where a brane is the boundary of a five-dimensional manifold. Now we will obtain a closed system of equations for a brane which is a junction between two volume spaces with the cosmological constants Λ_1 and Λ_2 , respectively, and without imposing Z_2 symmetry. Brane-world cosmologies without Z_2 symmetry of reflection were a subject of thorough study, for example, in [14,15,8,5]. Our study will be slightly more complete in the sense that we retain the curvature term in the action on the brane, i.e., in our case, $m \neq 0$. The braneworld model without Z_2 symmetry was studied in [16] in the case of $m \neq 0$ and $\Lambda_1 = \Lambda_2$.

Equation (4) now is modified as follows:

$$m^2 G_{ab} + \sigma h_{ab} = \epsilon M^3 (S_{ab}^{(1)} + S_{ab}^{(2)}) + \tau_{ab}, \quad (19)$$

where the tensors $S_{ab}^{(1)}$ and $S_{ab}^{(2)}$ are expressed through the extrinsic curvatures of the brane in two volumes, respectively. (We choose the normals n_1^a and n_2^a with respect to which we calculate the extrinsic curvatures to be the inner normals in the corresponding volumes.) We introduce the sum $S_{ab} = S_{ab}^{(1)} + S_{ab}^{(2)}$ so that Eq. (19) has the same form as Eq. (4).

Equation (8) is now satisfied by $S_{ab}^{(i)}$ in the place of S_{ab} and Λ_i in the place of Λ for $i = 1, 2$. Introducing also the difference $Q_{ab} = S_{ab}^{(1)} - S_{ab}^{(2)}$, we easily obtain the following closed system of gravitational equations on the brane:

$$\epsilon(R - \Lambda_1 - \Lambda_2) + \frac{1}{4} \left(S_{ab} S^{ab} - \frac{1}{3} S^2 \right) + \frac{1}{4} \left(Q_{ab} Q^{ab} - \frac{1}{3} Q^2 \right) = 0, \quad (20)$$

$$S_{ab} Q^{ab} - \frac{1}{3} S Q = 2\epsilon (\Lambda_1 - \Lambda_2), \quad (21)$$

$$D_a Q^a_b = 0, \quad (22)$$

where $Q = Q_{ab} h^{ab}$, and S_{ab} is formally given by the same equation as (9):

$$S_{ab} = \frac{\epsilon}{M^3} \left(m^2 G_{ab} + \sigma h_{ab} - \tau_{ab} \right). \quad (23)$$

Equation (22) follows from relation (5). This system of equations is to be solved for the metric and matter fields and for the symmetric tensor field Q_{ab} on the brane. Imposing Z_2 symmetry corresponds to setting $Q_{ab} = 0$ and $\Lambda_1 = \Lambda_2$ and is equivalent to the case of brane as a boundary (edge) considered above and described by Eq. (10) (these two cases are related by rescaling the constant M^3 by a factor of two).

We consider briefly the cosmological implications of system (20)–(23). Of course, even with the homogeneous and isotropic cosmological setting for the ordinary matter and gravity, the symmetric tensor Q_{ab} , in general, need not have homogeneous and isotropic form. However, here, for simplicity, we set it to be homogeneous and isotropic so that

$$Q^0_0 = \beta(t), \quad Q^\mu_\nu = \delta^\mu_\nu q(t), \quad \mu, \nu = 1, 2, 3. \quad (24)$$

Then Eqs. (21) and (22) easily yield the condition

$$2m^2 \beta \left(H^2 + \frac{\kappa}{a^2} - \frac{\rho + \sigma}{3m^2} \right) = M^3 \left(\Lambda_1 - \Lambda_2 + \frac{E}{a^4} \right), \quad (25)$$

where E is the integration constant. Now, using the conservation equation (22), we have

$$a^3 \dot{a} \left(Q_{ab} Q^{ab} - \frac{1}{3} Q^2 \right) = -\frac{1}{3} \frac{d}{dt} \left(a^4 \beta^2 \right). \quad (26)$$

Finally, a procedure quite similar to that which led us to Eq. (17) gives a partial generalization of Eq. (17) to the case where Z_2 symmetry is not imposed:

$$m^4 \left(H^2 + \frac{\kappa}{a^2} - \frac{\rho + \sigma}{3m^2} \right)^2 = 4\epsilon M^6 \left(H^2 + \frac{\kappa}{a^2} - \frac{\Lambda_1 + \Lambda_2}{12} - \frac{C}{a^4} \right) - \frac{M^{12}}{36m^4} \left[\frac{\Lambda_1 - \Lambda_2 + E/a^4}{H^2 + \kappa/a^2 - (\rho + \sigma)/3m^2} \right]^2, \quad (27)$$

where we used Eq. (25) to eliminate β . A similar equation was derived in [16] by the standard method of embedding for the partial case $\epsilon = 1$ and $\Lambda_1 = \Lambda_2$. The case of Z_2 symmetry corresponds here to setting $\Lambda_1 = \Lambda_2$ and $E = 0$. For $m = 0$, this equation reduces to a generalization of (18),

$$H^2 + \frac{\kappa}{a^2} = \frac{\Lambda_1 + \Lambda_2}{12} + \frac{C}{a^4} + \epsilon \frac{(\rho + \sigma)^2}{36M^6} + \epsilon \frac{M^6}{16} \left(\frac{\Lambda_1 - \Lambda_2 + E/a^4}{\rho + \sigma} \right)^2, \quad (28)$$

which in various forms appeared in [14] for the case of $\epsilon = 1$.

We conclude this work. The system of equations (20)–(23) is a closed system that describes the evolution on a brane. This system is all one has to solve in a theory in which there are no restrictions on solutions in the five-dimensional bulk. If there *are* such restrictions in the form of boundary and/or regularity conditions, then system (20)–(23) cannot be regarded as complete: only those its solutions are admissible which can be developed to solutions in the bulk satisfying the imposed regularity conditions. This illustrates the importance of the precise specification of boundary and/or regularity conditions in the bulk for a complete formulation of the braneworld theory: in the absence of such specification, *all* the information about the brane is contained in the closed system (20)–(23).

Finally, we note that system (20)–(23) admits solutions with flat vacuum branes without fine tuning the constants of the theory even in the case $\Lambda_1 = \Lambda_2$, although it requires the absence of Z_2 orbifold symmetry and, as a consequence, absence of the Lorentz symmetry of the bulk [17].

This work was supported in part by the INTAS grant for project No. 2000-334.

[1] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [[hep-ph/9905221](#)]; Phys. Rev. Lett. **83**, 4690 (1999) [[hep-th/9906064](#)].

- [2] P. Binétruy, C. Deffayet, U. Ellwanger, and D. Langlois, Phys. Lett. B **477**, 285 (2000) [[hep-th/9910219](#)].
- [3] K. Koyama and J. Soda, *Bulk gravitational field and cosmological perturbations on the brane*, Report No. KUCP0187 [[hep-th/0108003](#)].
- [4] R. Maartens, Phys. Rev. D **62**, 084023 (2000) [[hep-th/0004166](#)]; C. Gordon and R. Maartens, Phys. Rev. D **63**, 044022 (2001) [[hep-th/0009010](#)]; D. Langlois, R. Maartens, M. Sasaki, and D. Wands, Phys. Rev. D **63**, 084009 (2001) [[hep-th/0012044](#)].
- [5] G. Kofinas, JHEP **0108**, 034 (2001) [[hep-th/0108013](#)].
- [6] V. Sahni and Yu. V. Shtanov, *New vistas in braneworld cosmology*, Honorable Mention in the 2002 Essay Competition of the Gravity Research Foundation, [gr-qc/0205111](#).
- [7] R. M. Wald, *General Relativity*, The University of Chicago Press, Chicago (1984).
- [8] H. Collins and B. Holdom, Phys. Rev. D **62**, 105009 (2000) [[hep-ph/0003173](#)].
- [9] G. Dvali, G. Gabadadze, and M. Porrati, Phys. Lett. B **485**, 208 (2000) [[hep-th/0005016](#)].
- [10] Yu. V. Shtanov, *On brane-world cosmology*, [hep-th/0005193](#).
- [11] P. Binétruy, C. Deffayet, and D. Langlois, Nucl. Phys. B **565**, 269 (2000) [[hep-th/9905012](#)].
- [12] T. Shiromizu, K. Maeda, and M. Sasaki, Phys. Rev. D **62**, 024012 (2000) [[hep-th/9910076](#)].
- [13] C. Deffayet, Phys. Lett. B **502**, 199 (2001) [[hep-th/0010186](#)].
- [14] P. Kraus, JHEP **9912**, 011 (1999) [[hep-th/9910149](#)]; D. Ida, JHEP **0009**, 014 (2000) [[gr-qc/9912002](#)]; H. Stoica, H. Tye, and I. Wasserman, Phys. Lett. B **482**, 205 (2000) [[hep-th/0004126](#)]; P. Bowcock, C. Charmousis, and R. Gregory, Class. Q. Grav. **17**, 4745 (2000) [[hep-th/0007177](#)]; A.-C. Davis, I. Vernon, S. C. Davis, and W. Perkins, Phys. Lett. B **504**, 254 (2001) [[hep-ph/0008132](#)]; B. Carter and J.-P. Uzan, Nucl. Phys. B **406**, 45 (2001) [[gr-qc/0101010](#)]; R. A. Battye, B. Carter, A. Mennim, and J.-P. Uzan, Phys. Rev. D **64**,

124007 (2001) [[hep-th/0105091](#)].

[15] N. Deruelle and T. Doležal, Phys. Rev. D **62**, 103502 (2000) [[gr-qc/0004021](#)]; W. B. Perkins, Phys. Lett. B **504**, 28 (2001) [[gr-qc/0010053](#)].

[16] N. J. Kim, H. W. Lee, and Y. S. Myung, Phys. Lett. B **504**, 323 (2001) [[hep-th/0101091](#)].

[17] Yu. V. Shtanov, *Flat vacuum branes without fine tuning*, [hep-th/0108211](#).